## Introduction to Birimian volcanology

Eudes S.N. Tegan<sup>1,\*</sup>, P.C. Hayman<sup>2</sup>, Inza Coulibaly<sup>1</sup>, Alain N. Kouamelan<sup>1</sup>

<sup>1</sup>Laboratoire de Géologie du Socle et de Métallogénie, UFR-STRM, Université FHB, 22 BP 582 Abidjan

22, Côte d'Ivoire; <sup>2</sup>Faculté des Sciences et de l'Ingénierie, Science de la terre et biologie, géologie et géochimie, Queensland University of Technology, 2 George St, Brisbane city QLD 4000, Australia

\*E-mail: ndrisati@gmail.com

Geodynamic models have listed ideas that the Birimian Gold Province is the result of subduction or rifting. The authors who have worked in this field have all shown the great diversity of formations that abounds the Birimian. In their studies, they just reported the presence of mafic volcaniclastics without giving more information about their origin. It is for this reason that this study was conducted to determine their origin. To do this, we used cartography, macroscopic and microscopic petrography and sedimentological analyses. All this has made it possible to define five lithofacies that are the coherent mafic and coherent felsic facies, the lava flow facies, the felsic volcaniclastic and volcanosediments and the mafic volcaniclastic facies. The facies of mafic volcanoclastics consist of pyroclastic fallout deposits, avalanche deposits, hyperconcentrated debris flow deposits and also river deposits. These formations would constitute the avalanche phase during the slide during the collapse stage of a volcanic structure. They are therefore epiclastics.

**Keywords**: Fettêkro, Birimian, avalanche debris, collapse, epiclastic, volcaniclastics

